MOTOMCO FARM & HOME

To Help With Indoor and Outdoor Norway Rat, Roof Rat and House Mice Control

Ready To Use

With Zinc Phosphide





This product may only be used inside and within 100 feet of man-made structures and for all burrow baiting

INTENDED FOR COMMERCIAL USE

PELLETED BAIT

KILLS RATS & MICE

ACTIVE INGREDIENT:

 Zinc Phosphide (CAS #1314-84-7)
 2%

 OTHER INGREDIENTS:
 98%

 TOTAL
 100%

Net Weight: 5 lbs (2.3 kg)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panels for First Aid and additional precautionary statements.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-854-2494 for emergency medical treatment information.

If you experience signs and symptoms such as nausea, abdominal pain, tightness in chest, or weakness, see a physician immediately. For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide information Center at 1.500-565-7378.

| If swallowed | Call a poison control center or doctor-immediately for treatment advice or transport the patient to the nearest hospital. Do not drink water. Do not administer anything by mouth or induce vomiting unless told to do so by the poison control center or doctor. |
|---------------------------|---|
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the phosphine-producing active, Zinc Phosphide. Probable mucosal damage may contraindicate the use of gastric lavage. For animals ingesting bait and/or showing poisoning signs, induce vomiting by using hydrogen peroxide. Sodium bicarbonate can be given orally to neutralize the stomach acidity. The stomach and intestinal tract can be evacuated, oxygen administered and cardiac and circulatory stimulants given.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, restrictions and precautions.

USE RESTRICTIONS:

Do not apply by any method, to any pest, or to any site not specified on this label.

- Only apply balt in locations out of reach of children, pets, domestic animals and nontarget wildlife. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the balt. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into balt compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas orner to vandalism.
- . Bait stations are mandatory for outdoor, above-ground use.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all persons out of the treated area during application.

This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash proceptacles, agricultural and public buildings, transport vehicles (ships, trains, airoutain).

AID MOTOMCO FARM & HOME



KILLS RATS & MICE

PELLETED BAIT

ACTIVE INGREDIENT:
Zinc Phosphide (CAS #1314-84-7) 2%
OTHER INGREDIENTS: 98%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds, fish and other wildlife. Dogs, cats and other animals might be poisoned if they leed upon animals that have eaten this bait. Do not contaminate water when disposing of equipment wash water or made.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or dispos

PESTICIDE STORAGE: Store only in original container lin a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be placed in trash

or delivered to an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke. Call your Local Waste Apency for any questions on proper disposal.

DISCLAIMER: To the extent permitted by applicable law, the seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent permitted by applicable law, the buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Net Weight: 5 lbs (2.3 kg)

EPA REG. NO.: 12455-131-3240

Available Exclusively From:

Madison, WI 53704 U.S.A. www.motomco.com

EPA EST. NO.: 12455-WI-1

The World Leader In Rodent Control Technology®

USE RESTRICTIONS (Continued from other panel)

docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

This product may be applied to active rodent burrows to control Norway rats, and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no locative Norway/roof rat burrows. Do not broadcast bait. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove bood trash, garbage, clutter and debris.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 4 pounds of bait.

APPLICATION DIRECTIONS:

SELECTION OF TREATMENT AREAS: After removing as much food, water and harborage as possible, determine dry, acid-free areas where rats and mice will most likely find and consume the ball. Generally these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rats and mice or their signs have been observed.

APPLICATION DIRECTIONS FOR RATS: Apply 1 to 2 teaspoon amounts of bait per placement. For outdoor use, place bait in active rat burrows or tamper-resistant bait stations. Maintain an uninterrupted supply of fresh bait until all signs of feeding have stooded. Do not treat the same area at less than 30 day intervals.

APPLCATION DIRECTIONS FOR MICE: Apply 1 teaspoon of bait per placement at 8 to 12 foot intervals. For outdoor use, place bait in tamper-resistant bait stations. Maintain an uninterrupted supply of fresh bait until all signs of feeding have stopped. Do not treat the same area at less than 30 day intervals. Collect and dispose of all dead animals.

FOLLOW-UP: Wearing waterproof glows, collect and dispose of dead, exposed animals. Dispose of lettover ball in accordance with disposal instructions. Use detergent and hot water to wash spoons used for application into burrows. Do not use spoon for mixing, holding or transferring lood or feed. To discourage enintlestation, limit sources of rodent food, water, and harborage as much as possible.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through the skin, or inhaled. Do not breathe dust. Avoid contact with eyes, skin, or clothing. Any person who retrieves careasses or unused bait following application of this product must wear gloves. Personal Protective Eduinment (PEI)

All handlers, including loaders and applicators must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Water-proof gloves

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product.

User Safety Recommendations

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
 Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

050112/04-12