

<u>MIXING</u>: Pour required amount into spray tank partially filled with water. Add KALIGREEN slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Add a spreader / sticker and /or other products last. Do not allow mixture to stand. Use within 24 hours as KALIGREEN may lose its activity if left in the tank.

<u>COMPATIBILITY</u>: KALIGREEN is compatible with most crop protection products and nutrients. However, do not mix it with highly acidic products as the effectiveness of KALIGREEN may be compromised. Use of a buffering agent for acidification of a tank mixture may also

decrease the effectiveness of KALIGREEN. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of DIRECTIONS: possible combinations that can be encountered, observe all precautions and limitations on the label of all products used in mixtures.

<u>TIMING</u>: Begin application at the first sign of disease. Repeat applications as new growth or infection occurs or as favorable disease conditions continue. Use in an Integrated Pest Management rotation program at 7-10 day intervals with other effective fungicides for resistance management. KALIGREEN may be applied up to the day before harvest (PHI=1 day) for all labeled crops except roses, turf, ornamentals and tobacco. The PHI for roses, turf and ornamentals is 4 hours; the PHI for tobacco is 10 days. KALIGREEN may also be applied post-harvest to remaining foliage to decrease overwintering inoculum.

For control of powdery mildew, use KALIGREEN as indicated on the following crops. Use the higher rate when disease pressure increases or a severe infection has occurred.

CROPS	RATE
Root and tuber vegetables, such as beet (garden, sugar), carrot	2 ¹ /2-3 lbs/A
Bulb vegetables, such as onions (dry bulb, green), garlic, leeks, shallots	2 ¹ /2-3 lbs/A
Leafy vegetables, such as endive, head and leaf lettuce, radicchio, spinach, Swiss chard	2 ¹ /2-3 lbs/A
Cole crops, such as broccoli, Brussels sprouts, cabbage, cauliflower	2 ¹ /2-3 lbs/A
Legume vegetables (succulent or dried), such as bean (snap, green), pea (edible-pod, field, green, snow)	21/2-3 lbs/A
Fruiting vegetables, such as eggplant, pepper (bell, chili, pimento), tomatillo, tomato	2 ¹ /2-3 lbs/A
Cucurbits, such as cucumber, cantaloupe, casaba, honeydew melon, pumpkin, summer squash (crookneck, zucchini), winter squash (butternut, acorn), watermelon	2 ¹ /2-5 lbs/A
Pome fruits, such as apple, pear, quince	2 ¹ /2-3 lbs/A
Stone fruits, such as apricot, cherry, nectarine, peach, plum, prune	2 ¹ /2-3 lbs/A
Grapes (wine, fresh market, raisin)	2 ¹ /2-5 lbs/A
Small fruits and berries, such as blackberry, blueberry, currant, loganberry, raspberry, strawberry	2 ¹ /2-3 lbs/A
Subtropical fruits, such as banana, date, mango, papaya, pineapple	2 ¹ /2-3 lbs/A
Cereal grains, such as barley, wheat	2 ¹ /2-3 lbs/A
Herbs and spices, such as basil, chives, coriander (cilantro), dill, oregano, parsley, thyme	2 ¹ /2-3 lbs/A
Hops	2 ¹ /2-5 lbs/A
Additional crops, mint (peppermint and spearmint), tobacco	2 ¹ /2-3 lbs/A
Ornamentals-field and greenhouse, such as bare root, container, bedding and	1-3 lbs/A
flowering stock, field grown cut flowers, vegetable transplants, nursery and landscape, potted flowering, shade and flowering trees, woody ornamentals*	or 1-3 lbs/100 gal.water/A for hand held applications
Roses (field and greenhouse)	1-3 lbs/A
	or 1-3 lbs/100 gal.water/A for hand held applications
Turf (ornamental lawns, golf courses and sod farms)	2 ¹ /2-3 lbs/A
	or 3 lbs/100 gal.water/A for hand held applications

*NOTE TO USER : Unintentional consequences such crop injury may result due to environmental or growing conditions, manner of use or application. Because plant tolerance of pesticides varies as conditions vary, treat a few plants under conditions expected to be encountered and observe for plant damage prior to full-scale application to large numbers of plants.

STORAGE AND DISPOSAL	may cont
Do not contaminate water, food or feed by storage or disposal.	To t SEL
Pesticide Storage: Store in a cool, drt place in- accessible to children. Pesticide Disposal: Wastes resulting from the use	cher reas usec unde
of this product must be disposed of on site or at an approved waste disposal facility. <u>Container Disposal:</u> Nonrefillable container. Do not reuse or refill this container. Completely empty bag into	TO T SELI MEF EXP
equipment. Then dispose of empty bag, by recycling if available, or in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.	To th shal dam prod mus

WARRANTY AND CONDITIONS OF SALE SELLERS OFFER AND THE BUYER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY The directions for the use of this product are believed to be adequate and must be followed carefully. However, it is impossible to take into account all the variables and to eliminate risks associated with its use. Injury or damage may result because of conditions that are beyond the control of the SELLER.

To the extent consistent with applicable law, SELLER warrants that this product conforms to the chemical description on this label and is believed to be reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under conditions of normal use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW,

SELLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

To the extent consistent with applicable law, the SELLER shall not be liable for consequential, special or indirect damages resulting from the use or handling of the product. Any variation or exemption from this warranty must be in writing and signed by an authorized representative of the SELLER.