

Brandt™ Caribbean Blend

Microbial Inoculant

CONTAINS NON PLANT FOOD INGREDIENTS

<i>Bacillus amyloliquefaciens</i>	100,000,000 CFU per gram
<i>Bacillus azotofomans</i>	100,000,000 CFU per gram
<i>Bacillus coagulans</i>	100,000,000 CFU per gram
<i>Bacillus firmus</i>	100,000,000 CFU per gram
<i>Bacillus licheniformis</i>	100,000,000 CFU per gram
<i>Bacillus megaterium</i>	100,000,000 CFU per gram
<i>Bacillus pumilus</i>	100,000,000 CFU per gram
<i>Bacillus subtilis</i>	100,000,000 CFU per gram
<i>Geobacillus stearothermophilus</i>	100,000,000 CFU per gram
<i>Paenibacillus durum</i>	100,000,000 CFU per gram
<i>Paenibacillus polymyxa</i>	100,000,000 CFU per gram
<i>Pseudeomonas aureofaciens</i>	22,000,000 CFU per gram
<i>Pseudomonas fluorescens</i>	22,000,000 CFU per gram
<i>Pseudomonas putida</i>	22,000,000 CFU per gram
<i>Streptomyces coelicolor</i>	22,000,000 CFU per gram
<i>Streptomyces griseus</i>	22,000,000 CFU per gram
<i>Streptomyces lydicus</i>	22,000,000 CFU per gram
<i>Streptomyces nodosus</i>	22,000,000 CFU per gram
<i>Trichoderma hamatum</i>	22,000,000 CFU per gram
<i>Trichoderma harzianum</i>	22,000,000 CFU per gram
<i>Trichoderma reesei</i>	22,000,000 CFU per gram

55.00% Dextrose, 20.00% Sucrose, 3.00% Amino Acids, 3.00% Brewers Yeast, 3.00% Active Yeast, 2.00% Humic Acid (leonardite) 1.25% Kelp, 0.75% Hydrated Sodium Calcium Aluminosilicate (Drying Agent)

Rate Recommendations

Tank mix 250 - 500 grams of BRANDT CARIBBEAN BLEND per hectare (use higher rate when abiotic stress is high), 100 ml of silica (to enhance microbial proliferation) and 4 liters of a quality paraffinic oil such as Brandt Saf-T-Add (acts as a sticking agent to promote foliar microbial colonization). Apply in sufficient

Distributed by:
NEWPORT - FERSAN (JAMAICA) LTD.
Newport East, Kingston 15, Jamaica
Tel. (876) 967-5815
Fax (876) 948-3772
Email: info@newportfersan.com

Net Weight: 10 Kg

volume of water to soak foliage thoroughly, or to soak solution into the soil profile.

Mixing and Handling Instructions

Do not apply during the heat of the day. Apply early morning, evening or during overcast conditions.

Do not tank mix with pesticides, but may be mixed with synthetic and organic fertilizers.

When used in rotation, BRANDT CARIBBEAN BLEND is best applied 48-72 hours after pesticide application.

Store in cool, dry location away from heat sources, direct sunlight or moisture.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Eyes: Moderately irritating to eyes. Eye protection should be worn while handling. In case of eye contact, flush eyes thoroughly with clear water for at least 10 minutes. If irritation persists, consult a physician immediately. **Skin:** Skin exposure may cause mild irritation. In case of skin contact, flush skin thoroughly with clear water for at least 10 minutes. If irritation persists, consult a physician immediately. **Internally:** Do not take internally. In case of ingestion, do not induce vomiting. Consult a physician immediately.

Warranty: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use; but neither this warranty nor any other warranty of merchantability or fitness of a particular product expressed or implied, extends to the use of this product contrary to label conditions, or under conditions not reasonably foreseeable to the seller; and buyer assumes the risk of any such use.

Brandt is a trademark of Brandt Consolidated Inc.



2014-04
Lot No:

BRANDT™

Manufactured by:
Brandt Consolidated, Inc.
2935 South Koke Mill Road
Springfield, Illinois 62711 USA
www.brandt.co
(217) 547-5840